

LMU-16/95

June, 1995

# An Isospin Analysis of $CP$ Violation in $B_d \rightarrow D\pi, D^*\pi$ and $D\rho$

Zhi-zhong Xing<sup>1</sup>

*Sektion Physik, Theoretische Physik, Universität München,  
Theresienstrasse 37, D-80333 München, Germany*

## Abstract

By use of current experimental data, we carry out an isospin analysis of the weak decays  $B \rightarrow D\pi, D^*\pi$  and  $D\rho$ . It is found that only in  $B \rightarrow D\rho$  the strong phase shift of two different isospin amplitudes can be approximately neglected. We derive some useful relations between the  $CP$ -violating measurables and the weak and strong transition phases, and illustrate the different effects of final-state interactions on  $CP$  violation in  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$ .

---

<sup>1</sup>Electronic address: Xing@hep.physik.uni-muenchen.de

In the standard electroweak model,  $CP$  violation is naturally described by a non-trivial phase in the Cabibbo-Kobayashi-Maskawa (CKM) matrix. To test the consistency of the standard model, the most promising way is to study the phenomena of  $CP$  violation appearing in weak decays of neutral  $B$  mesons [1, 2]. Apart from a variety of  $B_d$  decays to  $CP$  eigenstates, some exclusive  $|\Delta B| = |\Delta C| = 1$  transitions to non- $CP$  eigenstates are also expected to have large  $CP$  asymmetries [3, 4]. In most of the previous predictions of branching ratios and  $CP$  violation for the latter type of decays, the final-state interactions were either ignored or injudiciously simplified. Hence an improvement of those naive and model-dependent calculations is desirable today, in order to yield reliable numerical results and confront them with the experiments in the near future.

In this note we shall follow a model-independent approach, i.e., the isospin analysis, to study  $CP$  violation in the decay modes  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$ . An obvious advantage is that the branching ratios for some of these decays have been measured [5], and a full reconstruction of the remaining modes is available at either existing or forthcoming  $B$ -meson facilities. By use of the isospin relations and current data, we obtain some constraints on the magnitudes of final-state interactions in the above decays. It is found that only in  $B \rightarrow D\rho$  the strong phase shift of two different isospin amplitudes plays an insignificant role. We derive the analytical relations between the  $CP$ -violating measurables and the weak and strong transition phases, and illustrate the different effects of final-state interactions on  $CP$  violation in  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$ .

Let us begin with the decay modes  $\bar{B}_d^0 \rightarrow D^+\pi^-$ ,  $\bar{B}_d^0 \rightarrow D^0\pi^0$ ,  $B_u^- \rightarrow D^0\pi^-$  and their  $CP$ -conjugate counterparts. The effective weak Hamiltonians responsible for these processes have the isospin structures  $|1, \mp 1\rangle$ . After some calculations we obtain the following isospin relations:

$$\begin{aligned} M_{+-} &= \langle D^+\pi^- | \mathcal{H} | \bar{B}_d^0 \rangle = V_{cb}V_{ud}^* (A_{3/2} + \sqrt{2}A_{1/2}) , \\ M_{00} &= \langle D^0\pi^0 | \mathcal{H} | \bar{B}_d^0 \rangle = V_{cb}V_{ud}^* (\sqrt{2}A_{3/2} - A_{1/2}) , \\ M_{0-} &= \langle D^0\pi^- | \mathcal{H} | B_u^- \rangle = V_{cb}V_{ud}^* (3A_{3/2}) ; \end{aligned} \quad (1a)$$

and

$$\begin{aligned} N_{-+} &= \langle D^-\pi^+ | \mathcal{H} | B_d^0 \rangle = V_{ud}V_{cb}^* (A_{3/2} + \sqrt{2}A_{1/2}) , \\ N_{00} &= \langle \bar{D}^0\pi^0 | \mathcal{H} | B_d^0 \rangle = V_{ud}V_{cb}^* (\sqrt{2}A_{3/2} - A_{1/2}) , \\ N_{0+} &= \langle \bar{D}^0\pi^+ | \mathcal{H} | B_u^+ \rangle = V_{ud}V_{cb}^* (3A_{3/2}) . \end{aligned} \quad (1b)$$

Here  $A_{3/2}$  and  $A_{1/2}$  correspond to the isospin 3/2 and 1/2 amplitudes, whose CKM matrix elements have been factored out and whose Clebsch-Gordan coefficients have been absorbed into the definitions of  $A_{3/2}$  and  $A_{1/2}$ . In obtaining eq. (1), we have assumed that there is no mixture of  $B \rightarrow D\pi$  with other channels. It is clear that the above transition amplitudes form two isospin triangles in the complex plane:

$$M_{+-} + \sqrt{2}M_{00} = M_{0-} , \quad N_{-+} + \sqrt{2}N_{00} = N_{0+} . \quad (2)$$

Of course  $|M_{+-}| = |N_{-+}|$ ,  $|M_{00}| = |N_{00}|$  and  $|M_{0-}| = |N_{0+}|$  can be directly determined from measuring the branching ratios of  $B \rightarrow D\pi$ . Then we are able to extract the unknown quantities  $A_{3/2}$ ,  $A_{1/2}$  and the strong phase shift between them by use of eq. (1) or (2). Denoting  $A_{3/2}/A_{1/2} = re^{i\delta}$ , we relate  $r$  and  $\delta$  to the measurables through the following formulas:

$$r = \frac{1}{\sqrt{3(R_{+-} + R_{00}) - 1}}, \quad \cos \delta = \frac{r}{2\sqrt{2}} [3(R_{+-} - 2R_{00}) + 1], \quad (3)$$

where  $R_{+-} = |M_{+-}/M_{0-}|^2$  and  $R_{00} = |M_{00}/M_{0-}|^2$  are two observables independent of the uncertainty of the CKM factors. Since eq. (3) results only from the isospin calculations, it is very useful for a model-independent analysis of the relevant experimental data.

Similarly we can find the same isospin relations as eqs. (1-3) for the decay modes  $\bar{B}_d^0 \rightarrow D^{*+}\pi^-$ ,  $\bar{B}_d^0 \rightarrow D^{*0}\pi^0$ ,  $B_u^- \rightarrow D^{*0}\pi^-$ ;  $\bar{B}_d^0 \rightarrow D^+\rho^-$ ,  $\bar{B}_d^0 \rightarrow D^0\rho^0$ ,  $B_u^- \rightarrow D^0\rho^-$ ; and their  $CP$ -conjugate processes. The existing data on the above channels can be found in ref. [5]. For the purpose of illustration, here we only calculate  $R_{+-}$  and  $R_{00}$  by taking the central values or upper bounds of the measured branching ratios. The phase space differences induced by the mass differences  $m_{D^0} - m_{D^-}$ ,  $m_{\pi^0} - m_{\pi^-}$  and  $m_{\rho^0} - m_{\rho^-}$  are negligible, so is the life time difference  $\tau_{B_d} - \tau_{B_u}$  [5]. Our results of  $R_{+-}$  and  $R_{00}$  are listed in Table 1. Accordingly the lower bounds of the isospin parameters  $r$  and  $\cos \delta$  can be determined, as also shown in Table 1, with the help of eq. (3). These results are consistent with those obtained from the maximum likelihood method in ref. [6].

| Decay modes            | $R_{+-}$ | $R_{00}$  | $r$      | $\cos \delta$ |
|------------------------|----------|-----------|----------|---------------|
| $B \rightarrow D\pi$   | 0.566    | $< 0.090$ | $> 1.02$ | $> 0.78$      |
| $B \rightarrow D^*\pi$ | 0.500    | $< 0.186$ | $> 0.97$ | $> 0.47$      |
| $B \rightarrow D\rho$  | 0.582    | $< 0.041$ | $> 1.07$ | $> 0.94$      |

Table 1: Estimates of the decay rate ratios ( $R_{+-}$  and  $R_{00}$ ) and the isospin parameters ( $r$  and  $\delta$ ) for the decay modes  $B \rightarrow D\pi$ ,  $D^*\pi$  and  $D\rho$ .

From Table 1 we observe that the magnitudes of  $A_{3/2}$  and  $A_{1/2}$  are comparable in all three cases. The current data on  $B \rightarrow D\rho$  imply that  $\cos \delta_{D\rho}$  is close to unity, i.e., there are not significant final-state interactions in this type of decays. In contrast, the effects of final-state interactions on  $B \rightarrow D\pi$  and in particular on  $B \rightarrow D^*\pi$  are non-negligible, unless the branching ratios of  $\bar{B}_d^0 \rightarrow D^0\pi$  and  $D^{*0}\pi^0$  are extremely suppressed. Since  $\cos \delta \leq 1$ , the lower bound of  $R_{00}$  can be found from eq. (3) by fixing the value of  $R_{+-}$ :

$$R_{00} \geq \frac{1}{2} \left( 1 - \sqrt{R_{+-}} \right)^2. \quad (4)$$

Specifically, we have  $R_{00}(D\pi) \geq 0.031$ ,  $R_{00}(D^*\pi) \geq 0.043$  and  $R_{00}(D\rho) \geq 0.028$ . In Fig. 1 we plot the allowed ranges of  $r$  and  $\cos \delta$  as the functions of  $R_{00}$ . One can see that the smaller  $\delta$  is, the smaller  $R_{00}$  will be. Among the three groups of decay modes under discussion,  $B \rightarrow D\rho$  should be the best candidate for testing the factorization approximation and studying the  $CP$  asymmetries. Note that the lower bounds of  $R_{00}$  obtained above may model-independently isolate the branching ratios of  $\bar{B}_d^0 \rightarrow D^0\pi^0, D^{*0}\pi^0$  and  $D^0\rho^0$ . By use of the central values of the available data [5], we get  $\text{Br}(\bar{B}_d^0 \rightarrow D^0\pi^0) \geq 1.6 \times 10^{-4}$ ,  $\text{Br}(\bar{B}_d^0 \rightarrow D^{*0}\pi^0) \geq 2.2 \times 10^{-4}$  and  $\text{Br}(\bar{B}_d^0 \rightarrow D^0\rho^0) \geq 3.8 \times 10^{-4}$ . It is expected that these three modes can soon be established in experiments.

We proceed to discuss  $CP$  violation in the decay modes  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$ , where every final state is common to both  $B_d^0$  and  $\bar{B}_d^0$  mesons. The  $CP$  asymmetry is induced by the interplay of decay and  $B_d^0 - \bar{B}_d^0$  mixing [1, 2]. Taking  $B_d \rightarrow D\pi$  for example, we define the  $CP$ -violating interference term via  $B_d^0 \rightarrow D^+\pi^-$  ( $D^0\pi^0$ ) versus  $\bar{B}_d^0 \rightarrow D^+\pi^-$  ( $D^0\pi^0$ ) as  $\xi_{+-}$  ( $\xi_{00}$ ); and that via  $B_d^0 \rightarrow D^-\pi^+$  ( $\bar{D}^0\pi^0$ ) versus  $\bar{B}_d^0 \rightarrow D^-\pi^+$  ( $\bar{D}^0\pi^0$ ) as  $\zeta_{-+}$  ( $\zeta_{00}$ ). Explicitly, these measurables are expressed as

$$\xi_{+-} = \text{Im} \left( \frac{V_{td}V_{tb}^*}{V_{tb}V_{td}^*} \cdot \frac{M_{+-}}{\tilde{M}_{+-}} \right), \quad \xi_{00} = \text{Im} \left( \frac{V_{td}V_{tb}^*}{V_{tb}V_{td}^*} \cdot \frac{M_{00}}{\tilde{M}_{00}} \right), \quad (5a)$$

$$\zeta_{-+} = \text{Im} \left( \frac{V_{tb}V_{td}^*}{V_{td}V_{tb}^*} \cdot \frac{N_{-+}}{\tilde{N}_{-+}} \right), \quad \zeta_{00} = \text{Im} \left( \frac{V_{tb}V_{td}^*}{V_{td}V_{tb}^*} \cdot \frac{N_{00}}{\tilde{N}_{00}} \right), \quad (5b)$$

where the decay amplitudes  $\tilde{M}$  and  $\tilde{N}$  are given by [7]

$$\begin{aligned} \tilde{M}_{+-} &= \langle D^+\pi^- | \mathcal{H} | B_d^0 \rangle = V_{cd}V_{ub}^* \left( \tilde{A}_{3/2} - \sqrt{2}\tilde{A}_{1/2} \right), \\ \tilde{M}_{00} &= \langle D^0\pi^0 | \mathcal{H} | B_d^0 \rangle = V_{cd}V_{ub}^* \left( \sqrt{2}\tilde{A}_{3/2} + \tilde{A}_{1/2} \right); \end{aligned} \quad (6a)$$

and

$$\begin{aligned} \tilde{N}_{-+} &= \langle D^-\pi^+ | \mathcal{H} | \bar{B}_d^0 \rangle = V_{ub}V_{cd}^* \left( \tilde{A}_{3/2} - \sqrt{2}\tilde{A}_{1/2} \right), \\ \tilde{N}_{00} &= \langle \bar{D}^0\pi^0 | \mathcal{H} | \bar{B}_d^0 \rangle = V_{ub}V_{cd}^* \left( \sqrt{2}\tilde{A}_{3/2} + \tilde{A}_{1/2} \right). \end{aligned} \quad (6b)$$

For simplicity, we denote  $\tilde{A}_{3/2}/\tilde{A}_{1/2} = \tilde{r}e^{i\delta}$ , where  $\delta$  is the same as the strong phase shift in eqs. (1) and (3). Subsequently we use the Wolfenstein parameters [8] and the angles of the unitarity triangle [5] to express the CKM matrix elements. To a good degree of accuracy, we have  $V_{ud} \approx V_{tb} \approx 1$ ,  $V_{cd} \approx -\lambda$ ,  $V_{cb} \approx A\lambda^2$ ,  $V_{ub} \approx A\lambda^3\sqrt{\rho^2 + \eta^2}e^{-i\gamma}$  and  $V_{td} \approx A\lambda^3\sqrt{(1-\rho)^2 + \eta^2}e^{-i\beta}$ . Note that the parameters  $(\beta, \gamma)$  are dependent upon  $(\rho, \eta)$  through  $\tan \beta = \eta/(1-\rho)$  and  $\tan \gamma = \eta/\rho$ . The  $CP$ -violating terms  $\xi_{+-}$  ( $\zeta_{-+}$ ) and  $\xi_{00}$  ( $\zeta_{00}$ ) turn out to be:

$$\xi_{+-} = \frac{(r\tilde{r} - 2) \sin(2\beta + \gamma) + \sqrt{2}r \sin[\delta - (2\beta + \gamma)] + \sqrt{2}\tilde{r} \sin[\delta + (2\beta + \gamma)]}{\lambda^2\sqrt{\rho^2 + \eta^2} h(\tilde{r}^2 + 2 - 2\sqrt{2}\tilde{r} \cos \delta)}, \quad (7a)$$

$$\xi_{00} = \frac{(2r\tilde{r} - 1) \sin(2\beta + \gamma) - \sqrt{2}r \sin[\delta - (2\beta + \gamma)] - \sqrt{2}\tilde{r} \sin[\delta + (2\beta + \gamma)]}{\lambda^2\sqrt{\rho^2 + \eta^2} h(2\tilde{r}^2 + 1 + 2\sqrt{2}\tilde{r} \cos \delta)}; \quad (7b)$$

and

$$\zeta_{-+} = \frac{(2 - r\tilde{r}) \sin(2\beta + \gamma) + \sqrt{2}r \sin[\delta + (2\beta + \gamma)] + \sqrt{2}\tilde{r} \sin[\delta - (2\beta + \gamma)]}{\lambda^2 \sqrt{\rho^2 + \eta^2} h (\tilde{r}^2 + 2 - 2\sqrt{2}\tilde{r} \cos \delta)}, \quad (8a)$$

$$\zeta_{00} = \frac{(1 - 2r\tilde{r}) \sin(2\beta + \gamma) - \sqrt{2}r \sin[\delta + (2\beta + \gamma)] - \sqrt{2}\tilde{r} \sin[\delta - (2\beta + \gamma)]}{\lambda^2 \sqrt{\rho^2 + \eta^2} h (2\tilde{r}^2 + 1 + 2\sqrt{2}\tilde{r} \cos \delta)}, \quad (8b)$$

where  $h$  denotes the ratio  $\tilde{A}_{1/2}/A_{1/2}$ . It should be noted that  $\zeta_{-+}$  ( $\zeta_{00}$ ) can be obtained from  $\xi_{+-}$  ( $\xi_{00}$ ) by the replacements  $\beta \rightarrow -\beta$  and  $\gamma \rightarrow -\gamma$ . From eqs. (7) and (8) one can see that in general the strong phase shift  $\delta$  enters the  $CP$  asymmetries and plays a significant role. Only when  $\delta$  is vanishingly small, its effect on  $\xi_{+-}$  ( $\zeta_{-+}$ ) and  $\xi_{00}$  ( $\zeta_{00}$ ) can be safely neglected. In this case, we have

$$\xi_{+-} = -\zeta_{-+} = \frac{r + \sqrt{2}}{\tilde{r} - \sqrt{2}} \cdot \frac{\sin(2\beta + \gamma)}{\lambda^2 \sqrt{\rho^2 + \eta^2} h}, \quad (9a)$$

$$\xi_{00} = -\zeta_{00} = \frac{\sqrt{2}r - 1}{\sqrt{2}\tilde{r} + 1} \cdot \frac{\sin(2\beta + \gamma)}{\lambda^2 \sqrt{\rho^2 + \eta^2} h}. \quad (9b)$$

Indeed the conditions  $\xi_{+-} = -\zeta_{-+}$  and  $\xi_{00} = -\zeta_{00}$  were injudiciously taken in most of the previous works (see, e.g., refs. [3, 4]). Considering the isospin results of  $r$  and  $\delta$  given in Table 1 and Fig. 1, we know that only  $\cos \delta_{D\rho} \approx 1$  is an acceptable approximation to the current data. Thus the  $CP$  asymmetries in  $B_d \rightarrow D\rho$  are dominated by the  $\sin(2\beta + \gamma)$  term of  $\xi_{+-}$  ( $\zeta_{-+}$ ) or  $\xi_{00}$  ( $\zeta_{00}$ ). In contrast, the previous numerical predictions of  $CP$  violation in  $B_d \rightarrow D\pi$  and  $D^*\pi$  are questionable.

Now let us illustrate the effects of nonvanishing  $\delta$  on  $CP$  asymmetries in  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$ . Typically we take  $\lambda \approx 0.22$ ,  $\rho \approx -0.07$  and  $\eta \approx 0.38$  [9], and this corresponds to  $\beta \approx 19.6^\circ$  and  $\gamma \approx 100.4^\circ$ . Fixing  $R_{+-}$ , we still use the constraints on  $R_{00}$  obtained in Table 1, eq. (4) and Fig. 1. Since there is not any experimental information on the magnitudes of  $\tilde{A}_{3/2}$  and  $\tilde{A}_{1/2}$ , here we make a likely but unjustified approximation:  $h \approx 1$  and  $\tilde{r} \approx r$ , just for the purpose of simplicity and illustration. The changes of  $\xi_{+-}, \zeta_{-+}$  and  $\xi_{00}, \zeta_{00}$  with the allowed values of  $\cos \delta$  are shown in Figs. 2 and 3 respectively. One can see that in  $B_d \rightarrow D\pi$  and  $D^*\pi$  the effects of  $\delta$  are significant, and the approximation  $\xi_{+-} = -\zeta_{-+}$  or  $\xi_{00} = -\zeta_{00}$  is invalid to a large extent. In comparison, the final-state interactions in  $B_d \rightarrow D\rho$  may be negligible, if the decay rate of  $\bar{B}_d^0 \rightarrow D^0 \rho^0$  is further suppressed to allow for  $\cos \delta_{D\rho} \rightarrow 1$ . Although the strong phase shift  $\delta$  plays a non-negligible role in each of the above channels, its effect can be well isolated after the measurements of  $\bar{B}_d^0 \rightarrow D^0 \pi^0, D^{*0} \pi^0$  and  $D^0 \rho^0$ . Thus the determination of  $CP$  violation in  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$  remains promising in the near future.

The experimental scenarios for observing  $CP$  violation in the exclusive  $|\Delta B| = |\Delta C| = 1$  transitions have been discussed in the literature (see, e.g., ref. [2]). The basic signal for

$CP$  violation between a channel (e.g.,  $B_d^0 \rightarrow D^+\pi^-$ ) and its  $CP$ -conjugate counterpart ( $\bar{B}_d^0 \rightarrow D^-\pi^+$ ) is a nonvanishing ratio of the difference to the sum of their decay rates. For either coherent or incoherent  $B_d^0\bar{B}_d^0$  decays to the non- $CP$  eigenstates under study, the  $CP$  asymmetries are always proportional to  $(\xi_{+-} - \zeta_{-+})$  or  $(\xi_{00} - \zeta_{00})$ , as shown in ref. [10]. Both the time-integrated and time-dependent measurements are available to establish the  $CP$ -violating signals in  $B_d \rightarrow D\pi, D^*\pi$  or  $D\rho$ , after  $10^{7-8}$   $B_d^0\bar{B}_d^0$  events have been accumulated.

I would like to thank Prof. H. Fritzsch for his constant encouragement. My gratitude goes also to Prof. G. Ecker and Prof. H. Pietschmann at the Institut für Theoretische Physik of Universität Wien, where this paper was written, for their warm hospitality. A stimulating discussion with Prof. D.D. Wu is greatly appreciated. Finally I am indebted to the Alexander von Humboldt-Stiftung for its financial support.

## References

- [1] A.B. Carter and A.I. Sanda, Phys. Rev. Lett. 45 (1980) 952; I.I. Bigi and A.I. Sanda, Nucl. Phys. B193 (1981) 85; B281 (1987) 41.
- [2] For a review, see: I.I. Bigi, V.A. Khoze, N.G. Uraltsev, and A.I. Sanda, in *CP Violation*, edited by C. Jarlskog (World Scientific, Singapore, 1989), p. 175.; I. Dunietz, in *B Decays*, edited by S. Stone (World Scientific, Singapore, 1992), p. 393.
- [3] I. Dunietz and J.L. Rosner, Phys. Rev. D34 (1986) 1404; D. Du, I. Dunietz and D.D. Wu, Phys. Rev. D34 (1986) 3414.
- [4] F. Buccella, F. Lombardi, G. Miele and P. Santorelli, Z. Phys. C59 (1993) 437; D. Du, X. Li and Z. Xiao, Phys. Rev. D51 (1995) 279.
- [5] Particle Data Group, L. Montanet et al., Phys. Rev. D50 (1994) 1173.
- [6] H. Yamamoto, preprint HUTP-94/A006 (hep-ph/9403255).
- [7] R.G. Sachs, preprint EFI-85-22 (1985); D. Du, I. Dunietz and D.D. Wu, in ref. [3].
- [8] L. Wolfenstein, Phys. Rev. Lett. 51 (1983) 1945; M. Kobayashi, Prog. Theor. Phys. 92 (1994) 287; Z.Z. Xing, Phys. Rev. D51 (1995) 3958.
- [9] A. Ali and D. London, Z. Phys. C65 (1995) 431.
- [10] H. Fritzsch, D.D. Wu, and Z.Z. Xing, Phys. Lett. B328 (1994) 477; Z.Z. Xing, Phys. Rev. D50 (1994) R2957.




Figure 1: Possible ranges of the isospin parameters  $r = |A_{3/2}/A_{1/2}|$  and  $\delta = \arg(A_{3/2}/A_{1/2})$  allowed by the current data on  $B \rightarrow D\pi, D^*\pi$  and  $D\rho$ .





Figure 2: Illustration of the changes of the  $CP$ -violating terms  $\xi_{+-}$  and  $\zeta_{-+}$  with the allowed values of the strong phase shift  $\delta$  in the decay modes  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$ .







Figure 3: Illustration of the changes of the  $CP$ -violating terms  $\xi_{00}$  and  $\zeta_{00}$  with the allowed values of the strong phase shift  $\delta$  in the decay modes  $B_d \rightarrow D\pi, D^*\pi$  and  $D\rho$ .